

Communication via entangled coherent quantum network

A. El Allati ^a, Y. Hassouni ^a N. Metwally^b

^aFaculté des Sciences Département de Physique, Laboratoire de Physique Théorique URAC 13, Université Mohammed V - Agdal. Av. Ibn Battouta, B.P. 1014, Rabat, Morocco

^bMathematics Department, College of Science, University of Bahrain, P.O. Box, 32038 Bahrain

Abstract

A quantum network is constructed via maximum entangled coherent states. The possibility of using this network to achieve communication between multi-participants is investigated. We showed that the probability of teleported unknown state successfully, depends on the size the used network. As the numbers of participants increases, the successful probability does not depend on the intensity of the field. The problem of implementing quantum teleportation protocol via a noise quantum network is discussed. We show one can send information perfectly with small values of the field intensity and larger values of the noise strength. The successful probability of this suggested protocol increases abruptly for larger values of the noise strength and gradually for small values. We show that for small size of the used quantum network, the fidelity of the teleported state decreases smoothly, while it decreases abruptly for larger size of network.

Keywords: Quantum teleportation; Quantum network; Maximally entangled coherent states.

1 Introduction

Quantum teleportation is a way to send unknown quantum state remotely between two users, where it is destroyed at the sending station and appears perfectly at the receiving station. Since the first quantum teleportation protocol was first introduced by Bennett et al [1], there are a lot of attentions have been given to develop this phenomenon in many different directions. Among these directions using quantum channels described by continuous variables [2, 3]. Some efforts have been done to extent the used quantum channel multipartities systems [4, 5, 6]. In these types of quantum teleportation protocols, one needs a larger numbers of participants collaborate together to transfer the unknown information between any two of them. This process is called information transfer remotely over a quantum network [7].

There are some efforts have been done to generate entangled quantum network. As an example, Nguyen [8] has constructed a quantum network consists of 2^N parties of coherent states and used it to implement quantum teleportation. Brougham et. al [9] have used a passive quantum networks with logical

bus topology to transfer information safely. Also, Ciccarello et. al [10] have proposed a physical model for the systematic generation of N-partite states for quantum networking. On the other hand the presence of noise is associated with the deterioration of performance for quantum information tasks. Therefore investigating the dynamics of quantum network and its usefulness to achieve quantum information protocols is very important [11, 12].

This leads us to the aim of the current work, where we construct a maximum entangled coherent network, i.e. all the participants share maximum entangled coherent states. The properties of these entangled coherent states have been investigated extensively in [13]. Figure (1) describes a quantum network consists of four members, Alice, Bob, Clair and David, to send unknown information between Alice and David by the assistance of Bob and Clair. This Network is generalized to includes m participants and we investigate the possibility of using this network to transfer unknown information between any two members by the co-operation of the others. We have treated the problem of remotely transfer information over noise quantum network, where it is assumed that during the construction of the network the travelling states from the source to their locations in the network subject to noise. This type of noise is equivalent to employing a half mirror for the noise channel [14].

This article is organized as follows. In Sec.2, a teleportation protocol is proposed to transfer unknown information between two distinct partners via quantum network consists of four members. The generalized of this protocol is discussed in Sec.2.2. Achieving quantum teleportation over a noise quantum network is the subject of Sec.3. Finally, Sec.4 is devoted to discuss our results.

2 Teleportation via perfect quantum network

Entangled coherent states have been found to be useful for quantum teleportation. They are defined as

$$|\alpha\rangle_{cat}^{\pm} = N_{\alpha}^{\pm}(|\alpha\rangle \pm |-\alpha\rangle), \quad (1)$$

where, $N_{\alpha}^{\pm} = [2 \pm 2e^{-2|\alpha|^2}]^{-\frac{1}{2}}$ is a normalization factor and $|\pm\alpha\rangle$ are two coherent states with an equal complex amplitude α defined as

$$|\pm\alpha\rangle = \exp(-\frac{|\alpha|^2}{2}) \sum_{n=0}^{\infty} \frac{(\pm\alpha)^n}{\sqrt{n!}} |n\rangle. \quad (2)$$

This type of entangled coherent states have been used to teleport a Schrödinger cat state [27]. The tripartite entangled coherent state of the form

$$|\phi^{\pm}\rangle = N_{\alpha}^{\pm}(|\sqrt{2}\alpha, \alpha, \alpha\rangle \pm |-\sqrt{2}\alpha, -\alpha, -\alpha\rangle), \quad (3)$$

$N_{\alpha} = [2(1 \pm e^{-8|\alpha|^2})]^{-\frac{1}{2}}$, is used to teleport two qubits entangled coherent state [15]. Recently, El-Allati et al [13] have introduced a teleportation protocol




Figure 1: The scheme for teleporting the superposition coherent state ρ_A , from Alice to David within a network with assistance of other participants via quantum channel ρ^- , each one holds a photodetector

to teleport a tripartites coherent state via entangled coherent state of four partites defined as:

$$|\psi^\pm\rangle = N^\pm (|2\alpha, \sqrt{2}\alpha, \alpha, \alpha\rangle \pm |-2\alpha, -\sqrt{2}\alpha, -\alpha, -\alpha\rangle), \quad (4)$$

where $N^\pm = \sqrt{2(1 \pm \exp[-16|\alpha|^2])}$. Also, this protocol has been generalized to teleport a coherent entangled state of m modes by using entangled coherent state consists of $m + 1$ modes [13]. This class of states behaves as maximum entangled state, MES for $\rho^- = |\psi^-\rangle\langle\psi^-|$, where the concurrence [16] $\mathcal{C} = 1$ and as partially entangled state, PES for $\rho^+ = |\psi^+\rangle\langle\psi^+|$.

Encouraged by the work of Nguyen [8], we employ this class of the MES and its generalized version, which is described in [13], to teleport unknown state over quantum network. We perform this idea and its generalization on next subsections.

2.1 Teleportation through a network of four participants

Assume that Alice is asked to send unknown state ρ_{A_1} to David, where

$$\rho_{A_1} = \frac{1}{N_A^2} (|\kappa_1|^2 |2\alpha\rangle\langle 2\alpha| + \kappa_1 \kappa_2^* |2\alpha\rangle\langle -2\alpha| + \kappa_2 \kappa_1^* |-2\alpha\rangle\langle 2\alpha| + |\kappa_2|^2 |-2\alpha\rangle\langle -2\alpha|), \quad (5)$$

$\kappa_{1,2}$ are unknown complex numbers, and $N_A = \sqrt{|\kappa_1|^2 + |\kappa_2|^2 + 2e^{-8|\alpha|^2} \text{Re}(\kappa_2^* \kappa_1)}$. For this aim, the users use quantum network consists of four maximum entan-

gled coherent state given by,

$$|\psi\rangle_{ABCD} = N^- \left(|2\alpha, \sqrt{2}\alpha, \alpha, \alpha\rangle_{ABCD} - |-2\alpha, -\sqrt{2}\alpha, -\alpha, -\alpha\rangle_{ABCD} \right), \quad (6)$$

where $N^- = \sqrt{2(1 - \exp[-16|\alpha|^2])}$. The total state of the system is given by $|\psi_s\rangle = |\psi\rangle_{A_1} \otimes |\psi\rangle_{ABCD}$, where A, B, C and D referee to Alice, Bob, Clair and David respectively. To implement this protocol the network's members achieve the following steps:

1. Alice mixes the unknown state, $\rho_{A_1} = |\psi\rangle_{A_1} \langle\psi|$ with the quantum channel $\rho_{ABCD} = |\psi\rangle_{ABCD} \langle\psi|$ by applying a series of operations defined by beam splitters and phase shifters as, [15, 21]

$$\rho_{A_1} \otimes \rho_{ABCD}^- \rightarrow R_{A_1 A} \rho_{A_1} \otimes \rho_{ABCD}^- R_{A_1 A}^* = \rho_{out}, \quad (7)$$

where $R_{ij}|\mu\rangle|\nu\rangle = |\frac{\mu+\nu}{\sqrt{2}}\rangle|\frac{\mu-\nu}{\sqrt{2}}\rangle$ [13] and the output state, ρ_{out} is,

$$\begin{aligned} \rho_{out} = & |\kappa_1|^2 |\psi_1\rangle \langle\psi_1| - |\kappa_1|^2 |\psi_1\rangle \langle\psi_2| - \kappa_1 \kappa_2^* |\psi_1\rangle \langle\psi_3| + \kappa_1 \kappa_2^* |\psi_1\rangle \langle\psi_4| \\ & + |\kappa_1|^2 |\psi_2\rangle \langle\psi_2| + |\kappa_1|^2 \kappa_2^* |\psi_2\rangle \langle\psi_2| - \kappa_1 \kappa_2^* |\psi_2\rangle \langle\psi_3| + \kappa_1 \kappa_2^* |\psi_2\rangle \langle\psi_4| \\ & - \kappa_2 \kappa_1^* |\psi_3\rangle \langle\psi_1| + \kappa_2 \kappa_1^* |\psi_3\rangle \langle\psi_2| + |\kappa_3|^2 |\psi_3\rangle \langle\psi_3| + |\kappa_3|^2 |\psi_3\rangle \langle\psi_4| \\ & - \kappa_2 \kappa_1^* |\psi_4\rangle \langle\psi_1| - \kappa_2 \kappa_1^* |\psi_4\rangle \langle\psi_2| - |\kappa_2|^2 |\psi_4\rangle \langle\psi_3| + |\kappa_2|^2 |\psi_4\rangle \langle\psi_4| \end{aligned} \quad (8)$$

with,

$$\begin{aligned} |\psi_1\rangle &= |2\sqrt{2}\alpha, 0, \sqrt{2}\alpha, \alpha\rangle_{A_1 ABCD}, \quad |\psi_2\rangle = |0, 2\sqrt{2}\alpha, -\sqrt{2}\alpha, -\alpha\rangle_{A_1 ABCD} \\ |\psi_3\rangle &= |0, -2\sqrt{2}\alpha, \sqrt{2}\alpha, \alpha\rangle_{A_1 ABCD}, \quad |\psi_4\rangle = |-2\sqrt{2}\alpha, 0, -\sqrt{2}\alpha, -\alpha\rangle_{A_1 ABCD}. \end{aligned} \quad (9)$$

2. Alice performs two photon number measurements on modes A_1 and A using two detectors D_{A_1} and D_A , (see Fig.1). Bob and Clair should carry out the local number measurement of modes B and C by their detectors D_B and D_C , respectively. There are two different possibilities due to Alice's operations:

- *Alice's measurements such that $n_{A_0} = 0, n_A > 0$*

The participants, Alice, Bob and Clair send their measurement outcomes to David via a public channel, where $n_A + n_B + n_C = \text{odd}$. In this case, the state at David's hand collapses into,

$$\rho'_D = \lambda_1 |-\alpha\rangle \langle-\alpha| - \lambda_2 |-\alpha\rangle \langle\alpha| - \lambda_3 |\alpha\rangle \langle-\alpha| + \lambda_4 |\alpha\rangle \langle\alpha|, \quad (10)$$

where $\lambda_1 = \frac{|\kappa_1|^2}{N_1}$, $\lambda_2 = \frac{\kappa_1 \kappa_2^*}{N_1} (-1)^{n_A + n_B + n_C}$, $\lambda_3 = \frac{\kappa_2 \kappa_1^*}{N_1} (-1)^{n_A + n_B + n_C}$, $\lambda_4 = \frac{|\kappa_2|^2}{N_1}$, and $N_1 = |\kappa_1|^2 + |\kappa_2|^2 - 2(-1)^{n_A + n_B + n_C} e^{-2|\alpha|^2} \text{Re}(\kappa_2^* \kappa_1)$ is the normalized factor.

Finally, David applies the operator $P(\pi)$ (phase shifter) to Eq.(10) to get the final state $\rho_D = P(\pi)\rho'_D P^*(\pi)$, which is exactly the state $\rho_0 = |\psi\rangle_0\langle\psi|$, with a probability of success in given by,

$$\mathcal{P}_1 = \frac{e^{-3|\alpha|^2}}{4 \sinh(8|\alpha|^2)} \{ \sinh(11|\alpha|^2) - \sinh(3|\alpha|^2) \}. \quad (11)$$

- *Alice's measurements such that $n_{A_1} > 0$, $n_{A_2} = 0$.*

In this case David's state collapse into,

$$\rho''_D = \lambda_1 |\alpha\rangle\langle\alpha| - \lambda_2 |\alpha\rangle\langle-\alpha| - \lambda_3 |-\alpha\rangle\langle\alpha| + \lambda_4 |-\alpha\rangle\langle-\alpha|, \quad (12)$$

where $\lambda_1 = \frac{|\kappa_1|^2}{N_1}$, $\lambda_2 = \frac{\kappa_1 \kappa_2^*}{N_1} (-1)^{n_{A_1} + n_B + n_C}$, $\lambda_3 = \frac{\kappa_1^* \kappa_2}{N_1} (-1)^{n_{A_1} + n_B + n_C}$, $\lambda_4 = \frac{|\kappa_2|^2}{N_1}$ and $N_1 = |\kappa_1|^2 + |\kappa_2|^2 - 2(-1)^{n_{A_1} + n_B + n_C} e^{-2|\alpha|^2} \text{Re}(\kappa_2^* \kappa_1)$ is the normalized factor.

However if $n_A + n_B + n_C$ is odd, then nothing should be done by David and the teleported state is obtained with a probability of success,

$$\mathcal{P}_2 = \frac{e^{-3|\alpha|^2}}{4 \sinh(8|\alpha|^2)} \{ \sinh(11|\alpha|^2) - \sinh(3|\alpha|^2) \}, \quad (13)$$

From Eqs.(11&13), the probability in both cases are equal, $\mathcal{P}_1 = \mathcal{P}_2$. Therefor the total probability of successful teleportation is given by,

$$\mathcal{P} = \frac{e^{-3|\alpha|^2}}{2 \sinh(8|\alpha|^2)} \{ \sinh(11|\alpha|^2) - \sinh(3|\alpha|^2) \}, \quad (14)$$

which tend to $\frac{1}{2}$ in the limit $|\alpha| \rightarrow 0$ and to $\frac{1}{2}$ in the limit $|\alpha| \rightarrow \infty$.

- At the end, David can makes some operators on the bases $|\alpha\rangle \rightarrow |2\alpha\rangle$ and $|-\alpha\rangle \rightarrow | -2\alpha\rangle$ using the modified beam splitter for producing exactly the state of Alice $\rho_D = \rho_A$.

Figure.(2), describes the behavior of the succusses probability, $\mathcal{P}$ of achieving the quantum teleportation protocol via a network consists of four members sharing a maximum entangled coherent state defined by (8). It is clear that, for small values of the mean photon number i.e. $|\alpha|^2 \in [0, 0.7]$ the probability decreases as $|\alpha|^2$ increases. However, the minimum value of $\mathcal{P}$ in the interval $\alpha \in [0, 0.7]$ is almost $\simeq 0.43$ and in the limit $\alpha \rightarrow 0$, the probability tends to 0.5. On the other hand, the probability of successes is independent of α for $|\alpha|^2 > 1$, where $\mathcal{P} = 0.5$. Comparing our results with that depicted in [8], we can see that constructing a quantum network by using a maximum entangled state defined by (3) is much better. It is clear that in the earlier study [8], $\mathcal{P} \rightarrow 0.25$ in the limit $\alpha \rightarrow 0$, while in the current investigation $\mathcal{P} \rightarrow 0.5$ as $\alpha \rightarrow 0$. The second reason, the probability of successful teleportation is independent from the intensity of the field, for $|\alpha|^2 < 0.7$, while in [8] for $|\alpha|^2 < 3$. Third reason although, $\mathcal{P}$ decreases in the interval $\alpha \in [0, 0.7]$, the minimum probability still much better than that depicted in the earlier work.




Figure 2: The probability of successful teleportation $\mathcal{P}$ as function of $|\alpha|^2$ by continue line and compared it in the same case to the probability of successful of Nguyen [8] which defined by dash line.

2.2 Teleportation through a network of m participants

These results can be generalized to $m + 1$ participants on the quantum network by using maximally entangled coherent states. For this aim, we assume that the network's users share maximally entangled coherent state of the form

$$|\Psi\rangle_{net}^- = N_{m+1} (|2^{\frac{m-1}{2}}\alpha\rangle_m \dots |2^{\frac{1}{2}}\alpha\rangle_2 |\alpha\rangle_1 |\alpha\rangle_0 - |-2^{\frac{m-1}{2}}\alpha\rangle_m \dots |-2^{\frac{1}{2}}\alpha\rangle_2 |-\alpha\rangle_1 |-\alpha\rangle_0), \quad (15)$$

with N_{m+1} is the normalization factor given by

$$N_{m+1} = [2(1 - e^{-2^{m+1}|\alpha|^2})]^{-\frac{1}{2}} \quad (16)$$

The entangled and separable properties of this class of states are investigated in [13]. Also, these states are employed to perform a teleportation between any number of parties. In this context, we use them to achieve quantum teleportation over a network. The suggested protocol is implemented as following.

1. Assume that the user m , who shares a maximum entangled state (15) with the remaining $m - 1$ members, co-operate to send the state $|\psi_0\rangle$ to any member of the network. In this case, the generator of this type of the MES, sends the mode m to the emitter and the modes $m - 1$ to remaining users. The total state of the system is defined by $\rho_s = \rho_{A_0} \cdot \rho_{net}$ where $\rho_{net} = |\psi\rangle_{net} \langle \psi_{net}|$.
2. The user m performs a sequence of local operations (7) on his own state and the state $|\psi_0\rangle$. Then by using the two detectors D_0 and D_m , the user m counts the photon numbers in modes 0 and m .




Figure 3: The probability of successful teleportation $\mathcal{P}$ as function of $|\alpha|^2$, where the dash, solid and dot curves for a networks consists of 2, 3, 4 participants respectively.

3. The other users carry out the local number measurement of modes $m-1, m-2, \dots, 1$ by the local detectors $D_{m-1}, D_{m-2}, \dots, D_1$, respectively. As a result of these measurements there are two possibilities: *The first possibility is obtained for $n_0 = 0, n_m > 0$*

In this case, the state $|\psi_0\rangle$ collapses at the receiver into,

$$\rho'_0 = \lambda_1 |-\alpha\rangle\langle-\alpha| - \lambda_2 |-\alpha\rangle\langle\alpha| - \lambda_3 |\alpha\rangle\langle-\alpha| + \lambda_4 |\alpha\rangle\langle\alpha|, \quad (17)$$

where $\lambda_1 = \frac{|\kappa_1|^2}{N_m}$, $\lambda_2 = \frac{\kappa_1 \kappa_2^*}{N_m} (-1)^{n_m + n_{m-1} \dots + n_1}$, $\lambda_3 = \frac{\kappa_2 \kappa_1^*}{N_m} (-1)^{n_m + n_{m-1} \dots + n_1}$, $\lambda_4 = \frac{|\kappa_2|^2}{N_m}$ and $N_m = |\kappa_1|^2 + |\kappa_2|^2 - 2(-1)^{n_m + n_{m-1} \dots + n_1} e^{-2|\alpha|^2} \text{Re}(\kappa_2^* \kappa_1)$ is the normalized factor.

The remaining participants send their classical results via a public channel to the receiver, where $n_m + n_{m-1} \dots + n_1 = \text{odd}$, who applies the operator $P(\pi)$ and a series of operations $R_{i,j}$ defined by modified beam splitter for replica the original state on the state (17) to get the state $\rho_0 = P(\pi) \rho'_0 P^*(\pi)$.

The second possibility: *the measures of the partners such that, $n_{m+1} > 0, n_m = 0$* , in this case, one gets the same probability of successes as the previous case. So the information is transformed with an average probability given by,

$$\mathcal{P} = \frac{e^{-(2^{m-1}-1)|\alpha|^2}}{2 \sinh(2^m |\alpha|^2)} \{ \sinh((3 \cdot 2^{m-1} - 1)|\alpha|^2) - \sinh((2^{m-1} - 1)|\alpha|^2) \}. \quad (18)$$

Fig.(3), describes the dynamics of the successful probability of teleportation (18) for different size of the quantum network. It is clear that, for $m = 2$, i.e

two partners co-operate to send the information to the third, the probability, $\mathcal{P}$ decreases smoothly in the interval of $|\alpha|^2 \in [0, 0.25]$ and then increases gradually. However as $|\alpha|^2$ increases the probability increases and reaches to its maximum value at $|\alpha|^2 \simeq 1.6$. As one increases the numbers of participants, $\mathcal{P}$ decreases abruptly in a small range of $|\alpha|^2$ and increases faster than that depicted for less numbers of partners. However the minimum value of the probability of successes increases as one decreases the size of the network. Therefore, one can conclude that, by increasing the number of partners in the quantum network, the probability of success teleportation is independent of $|\alpha|^2$. So, one can perform a quantum teleportation in the presences of low field intensity with probability 0.5, by increasing the size of quantum network. On the other hand this figure confirms that the used quantum channel (15) is much better than that used in the earlier work cited in [8].

3 Teleportation via noise quantum network

Decoherence represents the most obstacles in the context of handling information. It arises during the interaction of the systems with their environments [22], imperfect devices [23], noise channel [24, 25], the energy loss or the photon absorption [14, 27, 29] and etc. Therefore, it is very important to investigate the dynamics of information in the presence of noisy [26].

Investigating the properties of quantum channels consists of coherent states have received considerable attentions. As an example, Van Enk [27] has considered the decoherence of multi-dimensional entangled coherent states due to photon absorption losses, where he calculated how fast that entanglement decays and how much entanglement is left. The entanglement degradation of entangled states suffering from photon absorption losses is investigated in [28]. The dynamics of multi entangled coherent states and the possibility of using them to perform quantum teleportation are investigated in [13].

In this section, we assume that we have quantum network, QN consists of a multi-entangled coherent states. Assume that we have a source supplies the partners, with a maximum entangled coherent states given by (15). These entangled coherent states propagate from the source to the locations of the partners. Due to the interaction with the environment, the maximum entangled coherent states turn into partial entangled states, where its degree of entanglement depends on the strength of the noise. Let us consider that the source produces MECS defined by the density operator, ρ^- given by Eq.(15). This entangled state turns into a partial entangled state defined by,

$$\rho_{PE} = U_{AE} \otimes U_{BE} \rho^- U_{BE}^\dagger \otimes U_{AE}^\dagger, \quad (19)$$

where $U_{IE}|\alpha\rangle|0\rangle_E = |\sqrt{\eta}\alpha\rangle_I|\sqrt{1-\eta}\alpha\rangle_E$, $I = A$, or B and $|0\rangle_E$ referees to the environment state. This effect is equivalent to employing a half mirror for the noise channel [14]. In an explicit form, one can write the output density

operator ρ_{PE} as,

$$\begin{aligned}\rho_{PE} = & \frac{1}{N_\alpha} \left[|2\sqrt{\eta}\alpha, \sqrt{2}\sqrt{\eta}\alpha, \sqrt{\eta}\alpha, \sqrt{\eta}\alpha\rangle \langle 2\sqrt{\eta}\alpha, \sqrt{2}\sqrt{\eta}\alpha, \sqrt{\eta}\alpha, \sqrt{\eta}\alpha| \right. \\ & + | -2\sqrt{\eta}\alpha, -\sqrt{2}\sqrt{\eta}\alpha, -\sqrt{\eta}\alpha, -\sqrt{\eta}\alpha\rangle \langle -2\sqrt{\eta}\alpha, -\sqrt{2}\sqrt{\eta}\alpha, -\sqrt{\eta}\alpha, -\sqrt{\eta}\alpha| \\ & - e^{-8(1-\eta)|\alpha|^2} |2\sqrt{\eta}\alpha, \sqrt{2}\sqrt{\eta}\alpha, \sqrt{\eta}\alpha, \sqrt{\eta}\alpha\rangle \langle -2\sqrt{\eta}\alpha, -\sqrt{2}\sqrt{\eta}\alpha, -\sqrt{\eta}\alpha, -\sqrt{\eta}\alpha| \\ & \left. - e^{-8(1-\eta)|\alpha|^2} | -2\sqrt{\eta}\alpha, -\sqrt{2}\sqrt{\eta}\alpha, -\sqrt{\eta}\alpha, -\sqrt{\eta}\alpha\rangle \langle 2\sqrt{\eta}\alpha, \sqrt{2}\sqrt{\eta}\alpha, \sqrt{\eta}\alpha, \sqrt{\eta}\alpha| \right] \quad (20)\end{aligned}$$

where $N_\alpha = 2(1 - e^{-16|\alpha|^2})$ is the normalized factor (for more details see [13]). Assume that the aim of Alice is sending unknown state ρ_{A_1} (5) to Divad through the noise quantum network (20). The partners follow the same steps as described in Sec.(2.1) to fulfill this task. If the members Alice, Bob and clair who performed the number measurement send their outcomes through a classical channel to David such that, $n_{A_1} = 0$ and $n_{A_2} > 0$ and $n_A + n_B + n_C$ is odd, then the final state at Divad's hand is

$$\begin{aligned}\rho_D = & \lambda_1 | -\sqrt{\eta}\alpha\rangle \langle -\chi| \langle -\sqrt{\eta}\alpha| \langle -\chi| - \lambda_2 | -\sqrt{\eta}\alpha\rangle \langle -\chi| \langle \sqrt{\eta}\alpha| \langle \chi| \\ & - \lambda_3 | \sqrt{\eta}\alpha\rangle \langle \chi| \langle -\sqrt{\eta}\alpha| \langle -\chi| + \lambda_4 | \sqrt{\eta}\alpha\rangle \langle \chi| \langle \sqrt{\eta}\alpha| \langle \chi|, \quad (21)\end{aligned}$$

where,

$$\begin{aligned}|\chi\rangle &= |2\sqrt{\eta'}\alpha\rangle_E |\sqrt{2}\sqrt{\eta'}\alpha\rangle_E |\sqrt{\eta'}\alpha\rangle_E |\sqrt{\eta'}\alpha\rangle_E \\ \lambda_1 &= \frac{|\kappa_1|^2}{N_\chi}, \quad \lambda_2 = \frac{\kappa_1 \kappa_2^*}{N_\chi} (-1)^{n_{A_2} + n_B + n_C}, \quad \lambda_3 = \frac{\kappa_2 \kappa_1^*}{N_\chi} (-1)^{n_{A_2} + n_B + n_C}, \quad \lambda_4 = \frac{|\kappa_2|^2}{N_\chi} \\ N_\chi &= |\kappa_1|^2 + |\kappa_2|^2 + e^{-2|\eta\alpha|^2} e^{-16|\sqrt{\eta'}\alpha|^2} \text{Re}(\kappa_2^* \kappa_1), \quad \eta' = 1 - \eta. \quad (22)\end{aligned}$$




Figure 4: The total probability, $\mathcal{P}_t$ of successful quantum teleportation via noise quantum network (20). The solid, dash and dot curves are evaluate for $\eta = 0.02, 0.05, 0.1$ respectively.

The probability of finding odd number of photons in this case is given by,

$$\mathcal{P} = \frac{N_{\chi} e^{-11|\sqrt{\eta}\alpha|^2}}{N_{A_1}^2 N_-^2} \{ \sinh(11|\sqrt{\eta}\alpha|^2) - \sinh(3|\sqrt{\eta}\alpha|^2) \}, \quad (23)$$

where, N_{A_1} is the normalization factor of the unknown state Eq.(5), and N_- is the normalization of quantum channel state (6).

In Fig.(4), the dynamics of the total successful probability, $\mathcal{P}_t = 2\mathcal{P}$ is investigated for different values of the noise strength, η . It is clear that, as one increases η , the probability, $\mathcal{P}_t$ increases faster while it increases gradually for small values of η . For larger values of $|\alpha|^2$, $\mathcal{P}_t \rightarrow \frac{1}{2}$, namely the probability of successes is independent of noise. However for small values of the field intensity, one can increase the probability of successes by increasing the noise strength. Therefore to send a coded information through a noise cavity with a large probability either one reduces the intensity of the field and increases the noise strength or increases the field intensity and reduces the noise strength.

The fidelity, $\mathcal{F}$ of the teleported state (5), via a noise quantum network defined by (20), is given by [31, 32],

$$\mathcal{F} = \text{tr}\{\tilde{\rho}_D \rho_{A_1}\} = \frac{f N_H + 1}{N_H + 1}, \quad (24)$$

where,

$$f = \frac{(1 - e^{-16\eta|\alpha|^2})(1 + e^{-16(1-\eta)|\alpha|^2})}{2(1 - e^{-16|\alpha|^2})}, \quad (25)$$

is the fidelity of the quantum network (8) and N_H is dimension of the Hilbert space.




Figure 5: (a) The fidelity of the teleported state (5) by using the noise quantum network (8) (b) the same as (a) but for $\eta = 0.9, 0.7, 0.5, 0.3, 0.1$ from top to bottom curves.

Fig.(5a), describes the dynamics of the fidelity, $\mathcal{F}$ in a small interval of the field intensity, i.e. $|\alpha|^2 \in [0, 1]$. It is clear that, for small values of η and α , the fidelity of the teleported state increases very fast and it increases smoothly for larger value of α . Therefore to send information with a unit fidelity for low intensity field, one has to increase the noise strength. In Fig.(5b), the fidelity $\mathcal{F}$ is plotted for different values of the noise strength η . It is shown that, for large values of $\eta = 0.95$ (most upper curve), the fidelity decreases smoothly to reaches to its lower limit ($\mathcal{F} = \frac{2}{3}$). As one decreases η , the fidelity decreases faster and reach its lower limit at $|\alpha|^2 \simeq \frac{3}{2}$. However for small values of $\eta \in [0, 0.5]$ the fidelity increases gradually and reaches its upper limit for larger values of the field intensity. On the other hand, the fidelity is independent of the noise strength in the limit of $|\alpha|^2 \rightarrow \infty$ for $\eta > 0.5$, while for any value of $\eta \leq 0.5$, the fidelity approaches to the fidelity of classical teleportation, $\mathcal{F} = \frac{2}{3}$.

From the preceding discussions, one can notice that the advantage of this current teleportation protocol is the probability of successes is much better than that obtained by Enk et.al [28]. Also, for $\eta = \frac{1}{2}$, the fidelity, $\mathcal{F} = \frac{2}{3}$, while in the former work [28] ($\mathcal{F} = \frac{1}{2}$). On the other hand, for any value of $\alpha \in [0, 1]$, the fidelity $\mathcal{F}$ is maximum for any value of η .

Finally, let us assume that the sender and receiver are a member of noise quantum network consists of m participants. The members follow the same steps as described in Sec.(2.2), they shall end up their protocol with a final state at the receiver's hand with a fidelity given by

$$\mathcal{F}_n = \frac{1}{3} \left[\frac{(1 + e^{-2^m(1-\eta)|\alpha|^2})(1 - e^{-2^m\eta|\alpha|^2})}{(1 - e^{-2^m|\alpha|^2})} + 1 \right]. \quad (26)$$

In Fig.(6), we investigate the dynamics of the fidelity, $\mathcal{F}_n$ for different size of the network, where we assume two different values of the noise strength. In Fig.(6a), we set a larger value of the noise strength, $\eta = 0.9$ and different size of the quantum network. It is clear that, for small size ($m = 3$), the fidelity, $\mathcal{F}_n$ decreases smoothly and reaches its minimum bound for larger values of the field intensity. However for larger values of m , $\mathcal{F}_n$ decreases faster and reaches its minimum bound for smaller values of the field intensity. Fig.(6b), describes the dynamics of $\mathcal{F}_n$ for a smaller value of $\eta \in [0, 0.5]$, where for small values of m , the fidelity increases gradually and reaches its upper bound for larger values of $|\alpha|^2$. However for larger size of the network, i.e m is larger, the fidelity increases very fast and reaches its upper value for small values of $|\alpha|^2$.

4 Conclusion

A quantum network consists of a multi-participant is constructed by using maximum entangled coherent state. This network is used to send information between any two members, where the other members co-operate with the sender to achieve this aim with high probability of successes. We showed as the size of the network increases, i.e, the number of members are more, the probability of the successful teleportation does not depend on the intensity of the field. This




Figure 6: (a) The fidelity of the teleported state (5) by using the noise quantum network (8) (b) the same as (a) but for $\eta = 0.9$, and $m = 3, 4, 5, 6, 7$ from top to bottom curves.

type of entangled coherent state is much better than that used in the earlier work of Nguyen [8], where in his proposal, the probability of successes reaches its maximum value for larger values of the intensity of the field ($|\alpha|^2 \geq 3$), while for the current work ($|\alpha|^2 \geq 0.7$).

The possibility of using this network to perform communication between its members in the presences of noise is investigated, where we assume that the travelling maximum entangled state from the sources to the locations of the members subject to noisy. Our results show that the probability of achieving quantum teleportation successfully increases faster for small values of the noise strength and reaches its maximum value for small values of the field intensity. However for larger value of the noise strength, the successful probability increases gradually and needs a larger values of the field intensity to be maximum.

The effect of the noise strength and the field intensity on the dynamics of the fidelity of the teleported state is investigated. We have found out that, one can send unknown information between any two members of the network with a unit fidelity for small values of the field intensity and larger values of the noise strength. However the fidelity decreases as one decreases the noise strength in $(0.5, 1]$ and increases as one increases the noise strength in $[0, 0.5)$. On the other hand, the fidelity reaches the classical limit ($\frac{2}{3}$) if we set the noise strength ($= \frac{1}{2}$) and for larger value of the field intensity, the fidelity of teleported state is independent of the noise strength.

This protocol is generalized to be used between any two members of a network consists of (m) users, where there are $(m - 2)$ members should co-operate with the sender to send unknown state safely to the receiver. Also, the possibility of implementing this generalized protocol in the presence of noise is discussed. The fidelity of the teleported state decreases smoothly for small size

of the network and abruptly for larger size.

References

- [1] C. H. Bennett *et al.*, Phys. Rev. Lett. **70**, 1895 (1993).
- [2] L. Vaidmann, Phys. Rev. A **49**, 1473 (1994).
- [3] S. L. Braunstein and H. J. Kimble, Phys. Rev. Lett. **80**, 869 (1998).
- [4] A. Karlsson and M. Bourennane, Phys. Rev. A **58**, 4394 (1998)
- [5] G. Rigolin, Phys. Rev. A **71**, 032303 (2005).
- [6] L. D. Chuang, and C. Z. Liang, Commun. Theor. Phys. **47**, 464 (2007).
- [7] S. Perseguers, M. Lewenstein, A. A. Acin, J. I. Cirac , Nature Physics **6**, 539 (2010); J.E. Gough, M. R. James, H.I. Phys. Rev. A **81**, 023804 (2010); A. Casaccino, S. Lloyd, S. Mancini, S. Severini. Int. J. of Quant. Inf. vol. **7** 1417 (2009).
- [8] Nguyen Ba An, Phys. Rev. A **68**, 022321 (2003).
- [9] T. Brougham, G. M. Nikolopoulos, I. Jex, Phys. Rev. A **80**, 052325 (2009).
- [10] F. Ciccarello, M. Paternostro, S. Bose, D. E. Browne, G. M. Palma, M. Zarcone, Phys. Rev. A **82**, 030302(R) (2010).
- [11] M. Plenio, S. Huelga, New J. Phys. **10**, 113019 (2008).
- [12] S. Perseguers Phys. Rev. A **81** 012310 (2010).
- [13] A. El. Allati, N. Metwally and Y. Hassouni, Opt. Commu., doi:10.1016/j.optcom.2010.08.041 (2010).
- [14] S.J. van Enk and O. Hirota, Phys. Rev. A **64**, 022313 (2001).
- [15] X. Wang, Phys. Rev. A **64**, 022303 (2001).
- [16] S. Hill and W.K. Wootters, Phys. Rev. Lett. **78**, 5022 (1997); W.K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
- [17] K. Mattle *at al.*, Phys. Rev. Lett. **76**, 4656 (1996).
- [18] E. Schrödinger, *Naturwiss.* **14**, 664 (1926).
- [19] R. J. Glauber "Coherent and Incoherent States of the Radiation Field", *Physical Review* **131**, 2766 (1963).
- [20] E. Schrödinger; *Naturewissenschaften* **23**, 807 (1935).
- [21] R. Tahira, M. Ikram, H. Nha, M. Zubairy, Phys. Rev. A **79**, 023816 (2009).

- [22] G. M. Palma, K. -A. Suominen and A. K. Ekert, Proc. R. Soc. London **A452** 567 (1996).
- [23] B. Georgeot and D. L. Shepelyansky, Phys. Rev.E **62** 3504 (2000).
- [24] M. Sugahara and S. P.Kruchinin, Mod. Phys. Lett. B**15** 473 (2001).
- [25] N. Metwally and M. W. M. R. B. Wahiddin, Applied Mathematics & Information Sciences, **1**, 23-33 (2007); N. Metwally, quant-ph: arXiv:1007.1446 (2010).
- [26] J.-Hong An, M. Feng and W.-Min Zhang, Quantum Information and Computation **9**, 0317 (2009).
- [27] S. J. Van Enk, Phys. Rev. A **72**, 022308 (2005).
- [28] S.J. Van Enk, O. Hirota, Phys. Rev. A **71**, 062322 (2005).
- [29] H. F. Hofmann, T. Ide, T. Kobayashi and A. Furusawa, Phys. Rev. A **62**, 062304 (2000).
- [30] S. Popescu, Phys. Rev. Lett. **72**, 797 (1994).
- [31] M. Horodecki, P. Horodecki and R. Horodecki, Phys. Rev. A **60**, 1888 (1999).
- [32] S. Oh *et al.*, Phys. Rev. A **66**, 022316 (2002).